arXiv:1702.05111v1 [quant-ph] 16 Feb 2017

A note on the Exact Green function for a quantum system decorated by two or more impurities.

M.L. Glasser

Dpto de Física Teórica, Facultad de Ciencias
Universidad de Valladolid

Physics Department, Clarkson University,
Potsdam, NY 13699-5820
(USA)

Abstract

The exact Green function is constructed for a quantum system, with known Green function, which is decorated by two delta function impurities.It is shown that when two such impurities coincide they behave as a single singular potential with combined amplitude. The results are extended to N impurities.

Keywords: Quantum Green function, Delta function Potential

1 Introduction

The one dimensional harmonic oscillator or square well, for example, for which the energy -dependent Green function $G_0(x, x'; E)$ is known, have been taken for many years as solvable models for semi-conductor quantum wells[1]. Frequently delta function potentials are placed at various points to simulate defects or impurities. In the case of a single such potential, $V(x) = \lambda\delta(x - a)$, the Green function for the composite system is known to be[2]

$$G(x, x'; E) = G_0(x, x') + \lambda \frac{G_0(x, a; E)G_0(a, x')}{1 - \lambda G_0(a, a; E)} \tag{1}$$

In this note a corresponding formula is derived for the case $V(x) = \lambda\delta(x - a) + \mu\delta(x - b)$.

2 Calculation

We first note that the same argument can be used for the time dependent-, as well as the energy-dependent Green functions, so we shall omit the third argument and write simply $G(x, x')$.

Beginning with the Dyson equation, noting that $G_0(x, y) = G_0(y, x)$

$$G(x, x') = G_0(x, x') + \int G_0(x, y)V(y)G(y, x')dy, \tag{2}$$

where the integration extends over the system domain, one has the set of equations

$$G(x, x') = G_0(x, x') + \lambda G_0(x, a)G(a, x') + \mu G_0(x, b)G(b, x') \tag{3}$$

$$G(a, x') = G_0(a, x') + \lambda G_0(a, a)G(a, x') + \mu G_0(a, b)G(b, x'), \tag{4}$$

$$G(b, x') = G_0(b, x') + \lambda G_0(a, b)G(a, x') + \mu G_0(b, b)G(b, x'). \tag{5}$$

The linear equations (4) annd (5) are easily solved for $G(a, x')$ and $G(b, x')$:

$$G(a, x') = \frac{G_0(a, x') + \mu[G_0(b, x')G_0(a, b) - G_0(a, x')G_0(b, b)]}{D} \tag{6}$$

$$G(b, x') = \frac{G_0(b, x') + \lambda[G_0(a, x')G_0(a, b) - G_0(b, x')G_0(a, a)]}{D} \tag{7}$$

with

$$D = [1 - \lambda G_0(a, a)][1 - \mu G_0(b, b)] - \lambda\mu[G_0(a, b)]^2. \tag{8}$$

By inserting (6) and (7) into (3) we obtain the desired expression

$$G(x,x') = G_0(x,x')$$
$$+\frac{1}{D}\{\lambda G_0(x,a)G_0(a,x') + \mu G_0(x,b)G_0(b,x')$$
$$+\lambda\mu[G_0(x,a)\,(G_0(a,b)G_0(a,x') - G_0(b,b)G_0(b,x'))$$
$$+G_0(x,b)\,(G_0(a,b)G_0(a,x') - G_0(a,a)G_0(a,x'))]\}. \qquad (9)$$

3 Discussion

By setting λ to 0 (9) reduces to (1), proving this expression as well. The most salient feature of (9) is the denominator D whose zeros form the exact spectrum of the composite system. For example, when a and b coincide, D reduces to $1-(\lambda+\mu)G(a,a)$ and (9) reduces to (1) with λ replaced by the amplitude $\lambda+\mu$. I.e. the two impurities combine to form one with combined amplitude. This generalizes the result of Rinaldi and Fasssari[3], for two identical defects symmetrically placed with respect to the center of a harmonic oscillator.

Two further points can be made. Nothing in the derivation of (9) restricts it to the line. If we accept the standard definition $\delta(\vec{x}) = \Pi_{j=1}^{d}\delta(x_j)$, then (9), and its consequences, are valid for d-dimensional quantum systems. This has been proven function-theoretically for the three dimensional harmonic oscillator with two symmetrically placed identical impurities by Albeverio, Fassari and Rinaldi4].

A second observation is that D is simply the Cramer determinant for the pair of simultaneous linear equations (4) and (5). In the case of impurity potential

$$V(x) = \sum_{j=1}^{N} \lambda_j \delta(x-a_j) \qquad (10)$$

there will be N such equations and the determinant is easily evaluated. Thus, for $N=3$

$$D = \prod_{j=1}^{3}[1-\lambda_j G_0(a_j,a_j)] - \sum_{i<j}\lambda_i\lambda_j G_0(a_i,a_j)G_0(a_j,a_i)$$
$$+\lambda_1\lambda_2\lambda_3 G_0(a_1,a_2)G(a_2,a_3)G_0(a_3,a_1). \qquad (11):$$

This reduces to the $N=1$ and $N=2$ cases appropriately and shows that any two coinciding impurities coalesce as indicated above.

Finally, by letting N become infinite, the analogue of (11) might offer a new approach to Kronig-Penney-type systems for periodic or random unit cells.

Acknowledgement: The author thanks Prof. S. Fassari and Prof. L.M. Nieto for helpful comments and acknowledges the financial support of MINECO (Project MTM2014-57129-C2-1-P) and Junta de Castilla y Leon (VA057U16)).

References

[1] S. Albeverio, F. Gesztesy, R.Hoegh-Krohn and H. Holden, *Solvabe Models in Quantum Mechanics*, [AMS-Chelsea, Providence (2004)]

[2] M.L. Glasser and L.M. Nieto, *The energy level structures of a variety of one-dimensional confining potentials and the effects of a local singular perturbation*, Can. J. Physics **93**, 1-9 (2015).

[3] 3] S. Fassari and F. Rinaldi, *On the spectrum of the Schroedinger Hamiltonian of the one-dimensional harmonic oscillator perturbed by two identical attractive point interactions*, Reports on Mathematical Physics **69**, 353-370 (2012).

[4] S.Albeverio, S. Fassari and F. Rinaldi,*Spectral properties of a symmetric three-dimensional quantum dot with a pair of identical attaractive impurities symmetrically situated around the origin*, Nanosystems, Physics, Chemistry and Mathematics **7**,268-289 (2016).